
Finally, we can write down the following Feynman-rules (given in momentum space) for the construction of S-matrix elements in a Klein-Gordon field theory with Φ^4 interaction:

- Draw all possible connected, topologically distinct Feynman-diagrams including loops with n external legs. Ignore vacuum-to-vacuum graphs (such as the $(\Delta(0))^2$ part above).
- For each external scalar field write 1.
- For each vertex of four scalar fields write $-i\lambda/4!$.
- For each internal line write the Feynman propagator:

$$\lim_{\epsilon \rightarrow 0} \frac{i}{(p^2 - m^2 + i\epsilon)}$$

- For each internal momentum corresponding to a loop write $\int d^4k/(2\pi)^4$.
- The four-momentum is conserved at each vertex.

- Divide by the proper symmetry factors, i.e. the number of ways of interchanging components of the diagram without changing it (e.g., for the $\mathcal{O}(\lambda)$ loop diagram $S = 2/4!$ and for the tree-level diagram $S = 1/4!$.)

Note that the factor z does not appear as part of the Feynman-rules. It will be re-introduced when calculating loop diagrams as part of the renormalization procedure.

The last step: S-matrix elements \rightarrow cross sections

The probability of the scattering of two particles with four-momenta q_1, q_2 and masses m_1, m_2 into n particles with four-momenta p_1, p_2, \dots, p_n and masses m_3, m_4, \dots, m_{n+2} is given by the differential cross section, $d\sigma$, as follows

$$d\sigma = \frac{1}{2\sqrt{[(2q_1 q_2)^2 - m_1^2 m_2^2]}} \prod_{i=1}^n \left[\frac{d^3 p_i}{(2\pi)^3 2p_i^0} \right] \\ (2\pi)^4 \delta^4 \left(\sum_{i=1}^n p_i - q_1 - q_2 \right) \overline{\sum} |\mathcal{M}_{fi}|^2$$

where $\overline{\sum}$ denotes averaging (summing) over the initial (final) state degrees of freedom (e.g., spin and color).

The invariant matrix element \mathcal{M}_{fi} is connected to the interaction part of the S-matrix, $\mathcal{M}_{fi} = \langle p_1 \dots p_n | T | q_1 \dots q_m \rangle$, and can be constructed using the Feynman-rules of the underlying Quantum Field Theory.

For a $2 \rightarrow 2$ scattering process of two particles with momenta q_1, q_2 and masses m_1, m_2 into 2 particles with momenta p_1, p_2 and masses m_3, m_4 , the differential cross section for the particle with momentum p_1 being scattered into the solid angle $d\Omega = d \cos \theta d\phi$ reads

$$\frac{d\sigma}{d\Omega} = \frac{1}{64\pi^2} \frac{1}{\sqrt{[(s - m_1^2 - m_2^2)^2 - 4m_1^2 m_2^2]}} \overline{\sum} |\mathcal{M}_{fi}|^2(s, t)$$

where we introduced the Lorentz-invariant kinematical variables (*Mandelstam variables*)

$$s = (q_1 + q_2)^2 = (p_1 + p_2)^2 \quad t = (q_1 - p_1)^2 = (q_2 - p_2)^2$$

$$u = (q_1 - p_2)^2 = (q_2 - p_1)^2$$

with $t = m_1^2 + m_3^2 - 2q_1^0 p_1^0 + 2|\vec{q}_1||\vec{p}_1| \cos \theta$.

It is convenient to work in the center-of-mass frame, where $\vec{q}_1 = -\vec{q}_2$, $\vec{p}_1 = -\vec{p}_2$ and s denotes the center-of-mass energy squared. In the limit of massless particles all momenta $|\vec{p}_i|$, $|\vec{q}_i|$ and energies p_i^0 , q_i^0 are equal to $\sqrt{s}/2$. The total cross section is obtained by integrating over the solid angle


$$\sigma = \int d\Omega \frac{d\sigma}{d\Omega} = \int_{-1}^1 d\cos\theta \int_0^{2\pi} d\phi \frac{d\sigma}{d\Omega}$$

3.6 The Feynman-rules for QED

For any given action (Lagrangian) we can determine the Feynman-rules to write down the perturbative expansion of the Quantum Field Theory.

As Feynman-rules for QED, the Quantum Field Theory of electromagnetic interactions among charged fermions, one finds (see, e.g., Bjorken and Drell II):

- Draw all possible connected, topologically distinct Feynman-diagrams including loops with n external legs. Ignore vacuum-to-vacuum graphs.
- For each external photon with momentum k associate a polarization vector $\epsilon_\mu(k, \lambda)$, if it is ingoing, and $\epsilon_\mu^*(k, \lambda)$ for an outgoing photon.
- For each vertex of two fermions and a photon write



where Q_f denotes the charge of the fermion (leptons: $Q_f = 1$).

- For each external fermion and anti-fermion draw a line with arrow, where the direction of the fermion and anti-fermion lines are opposite to each other. For each external (anti-)fermion with momentum p and spin s , write $u(p, s)(\bar{v}(p, s))$ for lines entering the graph and for lines leaving the graph write $\bar{u}(p, s)(v(p, s))$:

e^- in initial state – ingoing electron line



e^- in final state – outgoing electron line




e^+ in initial state – outgoing electron line



e^+ in final state – ingoing electron line



- For each internal (virtual) fermion and anti-fermion with momentum k and mass m draw a line with arrow and associate a propagator:



An example: e^+e^- annihilation into 2 photons

There are two Feynman-diagrams contributing to this process and the corresponding matrix element reads:

$$\begin{aligned} \mathcal{M} = & \bar{v}(q_2, s_2)(-ie\gamma_\mu) \frac{i}{(\not{q}_1 - \not{k}_1 - m + i\epsilon)} (-ie\gamma_\nu) u(q_1, s_1) \\ & \epsilon^{\nu*}(k_1, \lambda_1) \epsilon^{\mu*}(k_2, \lambda_2) + \\ & + \bar{v}(q_2, s_2)(-ie\gamma_\nu) \frac{i}{(\not{q}_1 - \not{k}_2 - m + i\epsilon)} (-ie\gamma_\mu) u(q_1, s_1) \\ & \epsilon^{\nu*}(k_1, \lambda_1) \epsilon^{\mu*}(k_2, \lambda_2) \end{aligned}$$

From this we obtain the spin averaged(summed) matrix element squared

$$\begin{aligned} \overline{|\mathcal{M}|^2} &= e^4 \sum_{\lambda_1, \lambda_2} (\epsilon^{\nu*}(k_1, \lambda_1) \epsilon^\sigma(k_1, \lambda_1)) (\epsilon^{\mu*}(k_2, \lambda_2) \epsilon^\rho(k_2, \lambda_2)) \\ & \frac{1}{4} \sum_{s_1, s_2} \left| \frac{\bar{v}\gamma_\mu(\not{q}_1 - \not{k}_1 + m)\gamma_\nu u}{(q_1 - k_1)^2 - m^2} + \frac{\bar{v}\gamma_\nu(\not{q}_1 - \not{k}_2 + m)\gamma_\mu u}{(q_1 - k_2)^2 - m^2} \right|^2 = \\ & = e^4 g^{\mu\rho} g^{\nu\sigma} \frac{1}{4} \sum_{s_1, s_2} \left| \frac{\bar{v}\gamma_\mu(\not{q}_1 - \not{k}_1 + m)\gamma_\nu u}{(q_1 - k_1)^2 - m^2} + \frac{\bar{v}\gamma_\nu(\not{q}_1 - \not{k}_2 + m)\gamma_\mu u}{(q_1 - k_2)^2 - m^2} \right|^2 \end{aligned}$$

This is a general feature of calculating cross sections for processes involving external fermions, i.e. that one will encounter expressions of the form

$$\sum_{s_1, s_2} |\bar{v}(q_2, s_2) \Gamma^{\mu\nu} u(q_1, s_1)|^2$$

with, e.g., $\Gamma^{\mu\nu} = \gamma^\mu \not{k} \gamma^\nu$ or $\Gamma^{\mu\nu} = \gamma^\mu \gamma^\nu$. Writing the above expression in spinor components and using the spin sum for Dirac spinors one finds

$$\begin{aligned} & \sum_{s_1, s_2} |\bar{v}(q_2, s_2) \Gamma^{\mu\nu} u(q_1, s_1)|^2 = \\ & \sum_{s_1, s_2} [\bar{v}(q_2, s_2) \Gamma^{\mu\nu} u(q_1, s_1)] [\bar{u}(q_1, s_1) \bar{\Gamma}^{\rho\sigma} v(q_2, s_2)] = \\ & = \sum_{s_1, s_2} [v_d(q_2, s_2) \bar{v}_a(q_2, s_2) \Gamma_{ab}^{\mu\nu} u_b(q_1, s_1) \bar{u}_c(q_1, s_1) (\bar{\Gamma}^{\rho\sigma})_{cd}] = \\ & = (\not{q}_2 - m_2)_{da} \Gamma_{ab}^{\mu\nu} (\not{q}_1 + m_1)_{bc} (\bar{\Gamma}^{\rho\sigma})_{cd} = \\ & = \text{Tr}\{(\not{q}_2 - m_2) \Gamma^{\mu\nu} (\not{q}_1 + m_1) \bar{\Gamma}^{\rho\sigma}\} \end{aligned}$$

For each continuous string of fermion lines in a Feynman-diagram such a trace over γ matrices has to be calculated.

The traces of Dirac matrices in 4 dimensions read:

$$\text{Tr}\{\gamma^\mu\} = 0 \text{ and for all other traces of odd numbers of } \gamma\text{'s}$$

$$\text{Tr}\{\gamma^\mu \gamma^\nu\} = 4g^{\mu\nu}$$

$$\text{Tr}\{\gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma\} = 4[g^{\mu\nu} g^{\rho\sigma} + g^{\mu\sigma} g^{\nu\rho} - g^{\mu\rho} g^{\nu\sigma}]$$

After having worked out the traces in our example we find:

$$\begin{aligned} \overline{\sum} |\mathcal{M}|^2 = 2e^4 & \left[\frac{(q_1 k_2)}{(q_1 k_1)} + \frac{(q_1 k_1)}{(q_1 k_2)} + 2m^2 \left(\frac{1}{(q_1 k_1)} + \frac{1}{(q_1 k_2)} \right) + \right. \\ & \left. -m^4 \left(\frac{1}{(q_1 k_1)} + \frac{1}{(q_1 k_2)} \right)^2 \right] \end{aligned}$$

This yields the differential cross section in the center-of-mass-frame as follows

($\tau = 4m^2/s, \beta = \sqrt{1 - \tau}$):

$$\begin{aligned} \frac{d\sigma}{d\Omega} &= \frac{\alpha^2}{s} \frac{1}{\beta} \left[\frac{1 + \beta^2 \cos^2 \theta}{\tau + \beta^2 \sin^2 \theta} + \frac{2\tau}{\tau + \beta^2 \sin^2 \theta} - \frac{2\tau^2}{(\tau + \beta^2 \sin^2 \theta)^2} \right] \\ &\rightarrow \frac{\alpha^2}{s} \left[\frac{1 + \cos^2 \theta}{\sin^2 \theta} \right] \text{ for } \tau \rightarrow 0 \text{ and } \sin \theta \neq 0 \end{aligned}$$

Finally, the exciting part - the comparison with experiment:
see, e.g., M.Derrick et al., Phys.Rev.D34, 3286 (1986).